



早期緩和療護導論

張正雄

彰濱秀傳紀念醫院

主題

- What's early palliative care?
- Why they need early palliative care?
- What is the efficacy?
- Integration of palliative care into standard oncology
- Advances in early palliative care in Taiwan
- Barriers to early palliative care



What's palliative
care ?



Hospice care – 安寧療護



The modern **hospice** movement began in the UK in the **1960s**. **Cicely Saunders**, a 20th century British nurse and social worker, was responsible for the formation of the core tenets applied in hospices around the world through her experiences at St Luke's Home.



The concepts of “total pain”, including physical, spiritual, and psychological discomfort; the proper use of opioids for patients with physical pain; attention to the needs of “family members and friends who **provide care for the dying**”,

Palliative care – 緩和療護



The term **palliative care** (in the setting of treatment given with the goal of symptom relief) was probably first coined by the Canadian surgeon **Balfour Mount in 1974**.



Three main features were developed, namely, **multidimensional assessment and management** of severe physical and emotional distress; **interdisciplinary care** by multiple disciplines in addition to physicians and nurses; emphasis on caring not only the **patients** but also for their **families** .

Supportive care - 支持性照護




Supportive care emerged as a concept and care approach in the late 1980s, somewhat later than palliative care, but with a similar **focus on the individual patient** with cancer, the host, **not the tumour**.



A new medical discipline aiming to provide predominantly cancer patients with support for the management of “**treatment-related effects**”

Definition of palliative care in 2002 declared by WHO - 定義


- Palliative care is an approach that improves the **quality of life** of patients and their families facing the problem associated with life-threatening illness, through the prevention and relief of suffering by means of early identification and impeccable assessment and treatment of pain and other problems, **physical, psychosocial and spiritual**
- 促進生活品質 / 身心靈




Five Main Ideas 五個主要重點

- (I) QOL focused approach - 生活品質
- (II) Whole-human approach - 全人
- (III) Care that encompasses both the patient and those involved with the patient (particularly caregivers) - 全家
- (IV) Respect patient autonomy and choice- 自主選擇
- (V) Support people through frank and thoughtful discussions on difficult subjects - 全程

Palliative care classification models – *Time-based model* 時程模式



Provider-based model 服務提供者模式





Why they need
early palliative
care?



Advocacy for early palliative care –
2010, Temel *et al.* announced a clinical trial concerning “early palliative care” in the *NEJM*

The NEW ENGLAND JOURNAL of MEDICINE

ORIGINAL ARTICLE


Early Palliative Care for Patients with Metastatic Non–Small-Cell Lung Cancer

Jennifer S. Temel, M.D., Joseph A. Greer, Ph.D., Alona Muzikansky, M.A.,
Emily R. Gallagher, R.N., Sonal Admane, M.B., B.S., M.P.H.,
Vicki A. Jackson, M.D., M.P.H., Constance M. Dahlin, A.P.N.,
Craig D. Blinderman, M.D., Juliet Jacobsen, M.D., William F. Pirl, M.D., M.P.H.,
J. Andrew Billings, M.D., and Thomas J. Lynch, M.D.


APPROACH TO THE PATIENT WITH INCURABLE CANCER

Palliative care needs in oncology

癌症患者緩和療護的需求



緩和療護的內容和時間表




What is the efficacy?



Early, Integrated Palliative Care in Patients with Metastatic Lung Cancer

轉移性肺癌臨床試驗




Palliative Care Model


- ❖ Palliative care provided by physicians and nurse practitioners
- ❖ Visits occurred in the Cancer Center (medical oncology, radiation oncology or chemotherapy visits).
- ❖ Oncology and palliative care visits were done in tandem or simultaneously.
- ❖ Visits were not scripted or prescribed.
- ❖ If patients were admitted to the hospital, they were followed by the palliative care team

早期緩和治療對患者報告指標的影響


Variable	Standard Care (N=47)	Early Palliative Care (N=60)	Difference between Early Care and Standard Care (95% CI)	P Value†	Effect Size‡
FACT-L score	91.5±15.8	98.0±15.1	6.5 (0.5–12.4)	0.03	0.42
LCS score	19.3±4.2	21.0±3.9	1.7 (0.1–3.2)	0.04	0.41
TOI score	53.0±11.5	59.0±11.6	6.0 (1.5–10.4)	0.009	0.52



病人存活率與生活品質的改變




Mean Change in Quality-of-Life Scores from Baseline to 12 Weeks in the Two Study Groups.



Kaplan-Meier Estimates of Survival According to Study Group.

早期緩和治療的好處 —— ENABLE II trial




Primary endpoints: patient-reported quality of life (QoL), symptom intensity, and resource use

Secondary endpoint: mood

早期緩和治療的好處

Patient outcomes



Patients, No.

Intervention	143	108	69	59	48	27	145	109	73	62	48	28	140	102	72	60	47	26
Usual care	130	97	74	54	44	31	134	100	76	54	45	31	128	98	76	54	44	31




Fig 1. CONSORT diagram. *One patient randomly assigned to standard care was erroneously assigned to early palliative care at the time of random assignment and therefore is included in the early palliative care study group.

化學治療利用率




Fig 2. Chemotherapy use between study groups in entire sample ($n = 147$). Rates of chemotherapy use did not differ significantly between groups for participants who received no chemotherapy (standard care [SC], three of 73 [4.1%] v early palliative care [PC], six of 74 [8.1%]; $P = .49$); first line only (SC, 27 of 73 [37.0%] v early PC, 21 of 74 [28.4%]; $P = .30$); second line (SC, 22 of 73 [30.1%] v early PC, 21 of 74 [28.4%]; $P = .86$); third line (SC, 12 of 73 [16.4%] v early PC, 14 of 74 [18.9%]; $P = .83$); and fourth line or more (SC, nine of 73 [12.3%] v early PC, 12 of 74 [16.2%]; $P = .64$). Four participants had missing chemotherapy data because they transferred care to other institutions, reducing sample size from 151 to 147.

化學治療利用率




Fig 3. Administration of final regimen of (A) intravenous and (B) oral chemotherapy at end of life ($n = 129$). Within 60 days of death, a significantly greater percentage of patients were receiving intravenous chemotherapy as final regimen in standard-care (SC) group compared with early palliative care (PC) group (SC, 31 of 67 [46.3%] v early PC, 15 of 62 [24.2%]; $P = .01$). Finding remained similar within 30 days (SC, 16 of 67 [23.9%] v early PC, seven of 62 [11.3%]; $P = .07$) and 14 days of death (SC, seven of 67 [10.4%] v early PC, one of 62 [1.6%]; $P = .06$), although not quite meeting threshold for statistical significance. Percentages of patients receiving oral chemotherapy did not differ significantly between groups within each of three time frames (all P values ranging from .67 to $> .99$).

按類別劃分的生命末期成本

	Standard Care N=65	Early Palliative Care N=60	Cost Difference
Inpatient Visits % of patients Mean cost (SD)	46% \$12,665 (20,580)	38% \$9,555 (17,275)	\$3,110
Outpatient Visits % of patients Mean cost (SD)	80% \$1,415 (1,649)	77% \$1,683 (2,027)	\$268
Chemotherapy % of patients Mean cost (SD)	42% \$1,654 (1,654)	28% \$1,014 (1,913)	\$640
Hospice Services % of patients Mean cost (SD)	65% \$1,808 (2,117)	70% \$2,933 (4,011)	\$1,125

Early palliative care might have more beneficial effects on **quality of life and intensity of symptoms** among patients with advanced cancer than among those given usual or standard cancer care alone. The effects are of clinical relevance for patients at an advanced disease stage with limited prognosis, when further decline in quality of life is the rule.



Cochrane
Library

Cochrane Database of Systematic Reviews

Early palliative care for adults with advanced cancer (Review)

Haun MW, Estel S, Rücker G, Friederich HC, Villalobos M, Thomas M, Hartmann M

早期緩和治療的好處

Improved QOL,

Quality of end-of-life care,

Decreased rates of depression,

Illness understanding,

Patient satisfaction

Integration of palliative care into standard oncology care



ASCO guideline

- The guideline states that, “*Inpatients and outpatients with advanced cancer should receive dedicated palliative care services, **early in the disease course, concurrent with active treatment**”, and*
- Strongly recommends “*Integration of palliative care into standard oncology care*”.

傳統與早期緩和治療


Traditional palliative care



Early palliative care



Proposed model of optimal oncology palliative care provision, including integration across providers and settings



緩和療護成功 融入腫瘤臨床 實務的指標



Components of integration from seven randomised trials

	Jordhøy et al (2000) ⁴⁷	Temel et al (2010) ³	Zimmermann et al (2014) ⁵	Bakitas et al (2015) ⁴⁸	Maltoni et al (2016) ⁴⁹	Temel et al (2017) ⁵⁰	Grønvold et al (2017) ⁵¹
Clinical structure							
Palliative care inpatient consultation team	Y	Y	Y	Y	..	Y	Y
Palliative care outpatient clinic	Y	Y	Y	Y	Y	Y	Y
Community-based care or home palliative care	Y	..	Y
Clinical processes							
Multidisciplinary specialised palliative care team	Y	Y	Y	Y	Y	Y	Y
Routine symptom screening in the outpatient oncology clinic
Administration of systemic cancer therapy (eg, chemotherapy and targeted agents) possible in patients admitted to palliative care service	Y	Y	Y	Y	Y	Y	Y
Follow prespecified palliative care guidelines	Y	Y	Y	Y	Y
Early referral to palliative care	Y	Y	Y	Y	Y	Y	Y
Availability of clinical care pathways (automatic triggers) for palliative care referral
Palliative care team routinely involved in multidisciplinary tumour conference for patient case discussions	..	Y
Communication, cooperation, and coordination between palliative and oncology service	Y	Y	Y
Routine discussion of prognosis, advance care planning with goals of care	Y	Y	Y	Y	Y	Y	Y


ORIGINAL RESEARCH

Early Palliative Care for Oncology Patients: How APRNs Can Take the Lead

HEIDI MASON,^{1,2} DNP, ACNP-BC, MARY BETH DERUBEIS,² MSN, FNP-BC, and
BETH HESSELTINE,² MSN, FNP-C

Abstract

Background: Patients with cancer need expert and multidisciplinary care throughout the trajectory of their illness. Palliative care should be instituted early in the course of their disease. Early palliative care enables patients and their families to control physical, psychological, social, and spiritual symptoms of the disease. In our current health-care system, early palliative care is not being integrated due to a lack of education of providers and nurses, an infrastructure that does not support palliative medicine, and poor communication skills among practitioners. **Methods and Results:** The Palliative Care Quiz for Nursing (PCQN) completed by nurse practitioners at a large Midwest cancer center found that those nurse practitioners had a poor understanding of the basic precepts of palliative care. This is consistent with the current literature. **Conclusion:** Advanced practice nurses should be educated on the principles of palliative care, as they are perfectly situated to advance the integration of early palliative care in the oncology setting.



台灣早期緩和療護的進展

The fourth phase of
the national cancer
prevention program
(2019-23)-- Health
Promotion
Administration

- **6 Strategies:**
- **5 (2)** The importance of hospice and palliative care:
- Since 2018, the HPA promoted that cancer prevention and treatment institutions should establish early care standards and procedures for providing early palliative care for patients with advanced cancer.

癌症診療品質認證

“Accreditation standard 3.3 (since 2019)

Cancer prevention and treatment institutions should establish **palliative care standards and procedures** for patients with advanced cancer.




Figure6-20 Scatter map of hospitals that passed cancer treatment quality certification in 2019

秀傳彰濱緩和照護團隊

目的，原則與目標

目的：

減輕痛苦，促進舒適

維護病人自主控制權

原則：

針對痛苦和不適症狀提供解除方案

整合病人與家屬心理和靈性層面的照顧

目標：

協助病人及其家屬獲得罹病後最佳的生活品質

服務項目

提供緩和照護的團隊服務

提供疾病治療流程與預後的充分資訊

面對死亡威脅的心理輔導與價值觀的再澄清

促進病人與家屬，病家與原治療團隊間之互動與了解

瀕臨死亡前後提供家屬支持及生活模式重建，並轉介安寧團隊的延續照護。

緩和照護 收案原則

新診斷晚期癌症病人

身體上有較嚴重的疾病症狀或心理、
社會及靈性方面的照顧需求

病人與家屬對於疾病，治療目標與預
後的認知不足


病人對於治療的選擇與疾病惡化後的
處理模式，明顯有下決定上的困難

家屬與病人，病家與原治療團隊間對
於治療流程與目標有溝通上的問題

緩和照護團隊之 醫療團隊成員

- 癌症緩和專科醫師、藥師、心理師、營養師、社工員、個案管理師等。

收案流程與篩選量表



緩和照護團隊篩選量表	這件事對你的生活所造成的影響程度?	結果
1. 得了癌症之後，我每日的活動量是： <input type="checkbox"/> 正常 <input type="checkbox"/> 減少了一點 <input type="checkbox"/> 減少，但每日在床上的時間不會超過一半 <input type="checkbox"/> 大量減少，而且每日有超過一半的時間在床上 <input type="checkbox"/> 嚴重減少，而且一天大部分時間在床上	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	由心理師進行： 疲憊評估_____分 因疲憊進行收案： <input type="checkbox"/> 是 <input type="checkbox"/> 否
2 得了癌症之後，我感到沮喪、憂鬱、無望： <input type="checkbox"/> 完全不會 <input type="checkbox"/> 部分日子 <input type="checkbox"/> 很常 <input type="checkbox"/> 大部分日子 <input type="checkbox"/> 所有日子	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	簡式健康量表評估_____分，因心理評估後進行收案 <input type="checkbox"/> 是 <input type="checkbox"/> 否
3. 得了癌症之後，我可以做： <input type="checkbox"/> 所有我想做的事 <input type="checkbox"/> 大部分我想做的事 <input type="checkbox"/> 很多我想做的事 <input type="checkbox"/> 部分我想做的事 <input type="checkbox"/> 完全不能做我想做的事	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	由營養師評估 進行營養評估_____分 因營養問題進行收案： <input type="checkbox"/> 是 <input type="checkbox"/> 否
4. 得了癌症之後，我經驗到的疼痛程度是： <input type="checkbox"/> 沒有 <input type="checkbox"/> 輕微 <input type="checkbox"/> 中等 <input type="checkbox"/> 嚴重 <input type="checkbox"/> 無法忍受的痛/頻繁地疼痛	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	由醫師或癌症個案管理師評估： 進行疼痛評估_____分， 因疼痛進行收案： <input type="checkbox"/> 是 <input type="checkbox"/> 否
5. 得了癌症對我家造成很大的負擔： <input type="checkbox"/> 完全不會 <input type="checkbox"/> 有時 <input type="checkbox"/> 大部分的日子是如此 <input type="checkbox"/> 每天當中大部分的時間是如此 <input type="checkbox"/> 總是（每天且時時刻刻）	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	由社工師進行評估 進行經濟評估，因經濟進行收案：
6. 得了癌症對我和我的家庭之經濟狀況影響程度： <input type="checkbox"/> 完全不會 <input type="checkbox"/> 輕微 <input type="checkbox"/> 中等 <input type="checkbox"/> 嚴重 <input type="checkbox"/> 非常嚴重	<input type="checkbox"/> 完全不會 <input type="checkbox"/> 中度影響 <input type="checkbox"/> 嚴重影響	<input type="checkbox"/> 是 <input type="checkbox"/> 否



Barriers to Early Palliative Care

了解腫瘤專科醫師對緩和療護的看法以及阻礙其使用的障礙




FIGURE 1: Patient referrals to palliative care. Chart courtesy of Cardinal Health Specialty Solutions.




FIGURE 2: Early introduction of palliative care. Chart courtesy of Cardinal Health Specialty Solutions.

Understanding Oncologists' Perceptions About Palliative Care and the Barriers Preventing Its Use



FIGURE 3: Challenges to delivering effective palliative care. Chart courtesy of Cardinal Health Specialty Solutions.






FIGURE 4: Training and education about palliative care. Chart courtesy of Cardinal Health Specialty Solutions.

Conceptual model for timely palliative care



及時轉診緩和療護的障礙和促進因素



結論

- 患有無法治癒癌症的患者，在整個疾病過程中都需要緩和療護
- 越來越多強有力的 1 級證據，顯示緩和治療對患者和負責照顧家屬的益處
- 已發表關於該主題的隨機試驗，顯示跟主要治療整合可以帶來病人健康上的益處，但整合的內容、時間和方式尚未完全確定
- 緩和療護已成為國家癌症防治計劃的一部分

Q&A

